

# A new algorithm for discussing Gröbner bases with parameters

ANTONIO MONTES <sup>\*1</sup>

<sup>1</sup>*Dep. Matemàtica Aplicada 2,  
Universitat Politècnica de Catalunya, Spain,  
e-mail: montes@ma2.upc.es*

## Abstract

Let  $F$  be a set of polynomials in the variables  $\bar{x} = x_1, \dots, x_n$  with coefficients in  $R[\bar{a}]$ , where  $R$  is a UFD and  $\bar{a} = a_1, \dots, a_m$  a set of parameters. In this paper we present a new algorithm for discussing Gröbner bases with parameters. The algorithm obtains all the cases over the parameters leading to different reduced Gröbner basis, when the parameters in  $F$  are substituted in an extension field  $K$  of  $R$ . This new algorithm improves Weispfenning CGB algorithm, obtaining a reduced complete set of compatible and disjoint cases. A final improvement determines the minimal singular variety outside of which the Gröbner basis of the generic case specializes properly. These constructive methods provide a very satisfactory discussion and rich geometrical interpretation in the applications.

## 1. Introduction.

In many practical applications it is required to determine a Gröbner basis of an ideal of polynomials whose coefficients depend on some parameters. The main problem in this context is to obtain the distinct reduced Gröbner basis for all possible values of the parameters.

Examples of this situation can be found, for instance, in constructive algebraic geometry when determining conditions for a family of curves to have singular points (see subsection 11.2); in robotics, in order to determine conditions on the magnitudes of a robot configuration with certain degrees of freedom and to solve the inverse kinematics problem (see subsection 11.3); in the load-flow problem for a given electrical network (Montes, 1995, 1998), (see subsection 11.4); in automatic theorem proving, and so on.

<sup>\*</sup>Partially supported by GEN. CAT. 1999SGR00356 and DGES-MEC PB98-0933.

A direct approach to this problem can be derived applying the Comprehensive Gröbner Basis (CGB) algorithm of Weispfenning (1992) (see also Becker & Weispfenning (1993)). The goal of Weispfenning’s algorithm is to obtain a Comprehensive Gröbner basis that specializes for all possible values of the parameters. For this purpose it constructs a Gröbner system, i.e. a complete set of constructible sets over the parameters, in order to add new polynomials to the initial basis and achieve the CGB. Nevertheless, the Gröbner system derived from CGB is, in general, not simple enough for applications, as it generates more cases than necessary and leads to a complex discussion. In this paper, we present an improved algorithm, called DISPGB, that abandons the objective of obtaining a CGB and focus its interest on the essentially different reduced Gröbner basis (Gröbner system), simplifying the general discussion. A dichotomic tree discussion is carried out using quasi-canonical representations of specialization. A further improvement of the output (GENCASE algorithm) provides a minimal singular variety, outside of which the Gröbner basis of the generic case is specialization-invariant, and a reduced complete set of special cases inside the given minimal singular variety. This is an important point that is not obtained by Weispfenning’s CGB algorithm. Although the question about the algorithm independence of the singular variety remains open, its practical determination by the algorithm provides a satisfactory discussion leading to a rich geometrical interpretation in applications (see section 11). All the algorithms described in the paper are implemented in *Maple V*, release 6. †

In the case of linear systems, Sit (1992) gives an algorithm for the discussion using determinants. For this case, determinants provide the theoretical elements for defining the minimal singular variety. Our algorithm does not use determinants at all, but provides a polynomial in the parameters directly related to the value of the system determinant. Moreover, this “discriminant” polynomial is also obtained for general non-linear systems, where no analogue of the system determinant is known.

In order to state precisely the problem, let  $F = \{f_1, \dots, f_s\} \subseteq R[\bar{x}, \bar{a}]$  be a set of polynomials in the variables  $\bar{x} = x_1, \dots, x_n$  and the parameters  $\bar{a} = a_1, \dots, a_m$ , where  $R$  is a unique factorization domain (UFD). We can consider  $F$  as a polynomial set in the variables  $\bar{x}$  with coefficients in  $R[\bar{a}]$ .

The goal of DISPGB is to obtain the distinct reduced Gröbner basis for all possible values of the parameters. Under this perspective, a case is a set of polynomial equalities and inequalities over the parameters  $\bar{a}$  accepting the same expression for the reduced Gröbner basis. It would be desirable to distinguish cases only when their corresponding reduced Gröbner bases have different leading power product sets in the variables  $\bar{x}$ . This is a very demanding requirement. Nevertheless, using GENCASE, this objective is always reached for the generic

†The program and help libraries implementing the algorithms and a tutorial containing all the applications given in the paper with its corresponding time evaluation are available at author’s url <http://www-ma2.upc.es/~montes>. Identification JSC3091.

case, and usually we get very close to it for the special cases as it will become clear in the examples.

The problem of specialization of Gröbner bases has been actually studied by many authors. The general basic problem can be formulated in the following way. Let  $\mathcal{R}, \mathcal{R}'$  be Noetherian commutative rings with identity, and  $\sigma : \mathcal{R} \rightarrow \mathcal{R}'$  be a ring homomorphism. When does a Gröbner basis  $G$  of an ideal  $I \subseteq \mathcal{R}[\bar{x}]$  map to a Gröbner basis of  $\langle \sigma(I) \rangle \subset \mathcal{R}'[\bar{x}]$  under the natural extension  $\sigma : \mathcal{R}[\bar{x}] \rightarrow \mathcal{R}'[\bar{x}]$ ?

In Kalkbrener (1997) conditions obtained by different authors for this question are reviewed (see Gianni (1987); Kalkbrener (1987); Adams, Boyle (1992); Pauer (1992); Gräbe (1993); Assi (1994); Becker (1994)). Kalkbrener (1997) also proves a necessary and sufficient condition when  $R'$  is a field.

We restrict ourselves to the case where the original ring  $\mathcal{R}$  is a polynomial ring  $R[\bar{a}]$  over a unique factorization domain  $R$  and  $\mathcal{R}'$  some extension field  $K$  of the quotient field  $\text{Quot}(R)$ . The variables  $\bar{a}$  are considered as parameters. In this case  $R[\bar{a}]$  can be embedded in the quotient field  $R(\bar{a})$  and we do not need the general algorithm for computing Gröbner basis over rings (Möller, 1988), but only the classical (Buchberger, 1965, 1985; Gebauer, Möller, 1987; Gianni, Mora, 1987) algorithm over fields.

The structure of this paper is as follows:

Sections 3 and 4 are devoted to the study of how specialization acts on the pseudo-division and Gröbner basis algorithms. It is proved that they specialize to the corresponding algorithms in  $K[\bar{x}]$  except for a  $m - 1$  dimensional variety (hypersurface). Any hypersurface outside which Buchberger's algorithm specializes will be called a *singular variety*. It is not difficult to complete the algorithm in order to obtain a singular variety.

Nevertheless, the singular variety computed by Buchberger's algorithm is in general larger than is strictly necessary, and is algorithm depending. In order to minimize it we introduce a Generalized Gaussian Elimination algorithm (GGE). This algorithm produces a new basis specializing properly for any specialization and is a good input for Buchberger's algorithm. It is also directly useful in applications, because it is able to simplify problems. Section 5 is devoted to the study of GGE algorithm (Montes, 1999).

In section 6 we describe how specializations can be specified in a quasi-canonical form in order to obtain a dichotomic discussion, that is a central point of DISPGB.

We are now ready to deal with the general DISPGB algorithm. Section 7 provides a general description of DISPGB that tries to make it understandable. Section 8 is devoted to two essential sub-algorithms. Section 9 is devoted to the control flow sub-algorithms; the final theorem describing DISPGB is also given there.

In section 10 the algorithm GENCASE, designed to improve the output of DISPGB is described, and the theorem allowing to define and determine the unique *generic case* and the associated *minimal singular variety* is proved.

Finally some classical applications are given in the last section, showing both the elegance and power of DISPGB.

## 2. Some notations

In what follows we assume that  $\succ_{\bar{x}}$  is a given monomial order for the variables  $\bar{x}$ . We use the following notations. A power product in the variables  $\bar{x}$  is denoted  $\bar{x}^\alpha$ . A monomial is a product of a coefficient in the coefficient ring or field and a power product of the variables. The leading monomial, coefficient and power product are denoted:

$$\text{lm}(f, \succ_{\bar{x}}) = c \bar{x}^\alpha, \quad \text{lc}(f, \succ_{\bar{x}}) = c, \quad \text{lpp}(f, \succ_{\bar{x}}) = \bar{x}^\alpha.$$

We will also need a monomial order  $\succ_{\bar{a}}$  involving the parameters  $\bar{a}$  and the compatible elimination product order  $\succ_{\bar{x}\bar{a}}$  (see Bayer & Stillman (1987)). It is defined by:

For all  $\alpha, \gamma \in \mathbb{Z}_{\geq 0}^n$  and  $\beta, \delta \in \mathbb{Z}_{\geq 0}^m$

$$\bar{x}^\alpha \bar{a}^\beta \succ_{\bar{x}\bar{a}} \bar{x}^\gamma \bar{a}^\delta \iff \{ \bar{x}^\alpha \succ_{\bar{x}} \bar{x}^\gamma \quad \text{or} \quad (\bar{x}^\alpha = \bar{x}^\gamma \quad \text{and} \quad \bar{a}^\beta \succ_{\bar{a}} \bar{a}^\delta) \}. \quad (1)$$

In order to avoid denominators and unnecessary factors in  $S$ -polynomials we define them as follows:

$$S(f, g) = \frac{\Gamma \bar{x}^\gamma}{\text{lm}(f)} f - \frac{\Gamma \bar{x}^\gamma}{\text{lm}(g)} g \quad (2)$$

where  $\Gamma = \text{lcm}(\text{lc}(f), \text{lc}(g))$  and  $\bar{x}^\gamma = \text{lcm}(\text{lpp}(f), \text{lpp}(g))$ , with a slightly different normalization than in Weispfenning (1992).

For specializations coming from replacement of parameters, the corresponding ring homomorphism is:

$$\sigma : R[\bar{a}] \longrightarrow K \quad (3)$$

where  $K$  is any field extension of  $R$  and  $\sigma|_R = \text{Id}$ . The discussion is complete when we consider the algebraic closure  $\bar{K}$  of  $K$ . Providing a specialization is equivalent to choosing  $\bar{a}_0 \in K^m$ , and setting  $\sigma(\bar{a}) = \bar{a}_0$ . We will denote also by  $\sigma$  the natural extension  $\sigma' : R[\bar{a}][\bar{x}] \longrightarrow K[\bar{x}]$  which is the identity on  $\bar{x}$ .

We can identify  $\sigma_{\bar{a}_0}$  with  $\bar{a}_0$ , and extend  $\sigma$  for certain elements of  $R(\bar{a})$  (more precisely to the local ring  $R[\bar{a}]_{\ker(\sigma)}$ ). When we say that  $\sigma(f)$  makes sense for  $f \in R(\bar{a})[\bar{x}]$  it has to be understood that  $f \in R[\bar{a}]_{\ker(\sigma)}[\bar{x}]$ .

For obtaining  $k$ -quasi-canonical representations of the specification, as well as in other parts of the algorithms in this paper, we need to decompose polynomials into irreducible factors. We denote

$$\text{FACVAR}(W, \succ_{\bar{a}}) = \{q_1, \dots, q_s\}$$

the set of irreducible polynomials that are factors of the polynomials of  $W$ , normalized in a canonical form in order to be recognized when compared.

Definition (2) is clearly specialization-invariant when the leading coefficients of  $f$  and  $g$  are assumed different from zero.

The ideal generated by the set  $F$  in the polynomial ring  $\mathcal{R}$  will be denoted  $\langle F \rangle$  or explicitly  $\langle F \rangle \mathcal{R}$ , when some doubt about the polynomial ring can arise.

### 3. Pseudo-division algorithm and specialization

We now study pseudo-division algorithm (see Cox, Little & O'Shea (1992) ) under specialization and prove the following

**THEOREM 1:** *Let  $f, g_1, \dots, g_s \in R[\bar{a}][\bar{x}]$ ,  $W = \text{FACVAR}(\text{lc}(g_1), \dots, \text{lc}(g_s))$  and  $\sigma$  a specialization satisfying  $\sigma(w) \neq 0$ , for all  $w \in W$ , so that  $\sigma(\text{lc}(g_i)) \neq 0$ , for  $1 \leq i \leq s$ . Let  $q_1, \dots, q_s \in R(\bar{a})[\bar{x}]$  and  $r \in R(\bar{a})[\bar{x}]$  be the quotients and remainder of the pseudo-division of  $f$  by  $g_1, \dots, g_s$  in  $R(\bar{a})[\bar{x}]$ . Then  $\sigma(q_1), \dots, \sigma(q_s)$  and  $\sigma(r)$  make sense, and they are the corresponding quotients and remainders for the pseudo-division of  $\sigma(f)$  by  $\sigma(g_1), \dots, \sigma(g_s)$  in  $K[\bar{x}]$ .*

*Proof:* We prove the result by induction on the number of division steps. Call  $(p^{(j)}, q_1^{(j)}, \dots, q_s^{(j)})$  and  $(p'^{(k)}, q_1'^{(k)}, \dots, q_s'^{(k)})$  the partial (remainders, quotients) at steps  $j$  and  $k$  respectively of the divisions in  $R(\bar{a})[\bar{x}]$  and in  $K[\bar{x}]$ . We want to prove that for each  $j$  there is some  $k$  so that  $\sigma(p^{(j)}) = p'^{(k)}$  and  $\sigma(q_i^{(j)}) = q_i'^{(k)}$  for  $i = 1, \dots, s$ . It is true, by construction, for  $j = 0$  by picking  $k = 0$ . Suppose we are at step  $j$  in the division in  $R(\bar{a})[\bar{x}]$ , and let us assume the induction hypothesis. Let  $i_j$  be the first index for which  $\text{lpp}(g_{i_j})$  divides  $\text{lpp}(p^{(j)})$ . The new partial quotient and the new partial remainder will be

$$\begin{aligned} \psi_{i_j} &= \frac{\text{lm}(p^{(j)})}{\text{lm}(g_{i_j})} \\ p^{(j+1)} &= p^{(j)} - \psi_{i_j} g_{i_j}. \end{aligned} \tag{4}$$

Observe that by hypothesis,  $\sigma$  makes sense on  $\psi_{i_j}$  and  $p^{(j+1)}$ . So, in specializing, two cases can occur:

Case 1:  $\sigma(\text{lc}(p^{(j)})) = 0$ . In this case  $\sigma(\psi_{i_j}) = 0$  and  $\sigma(p^{(j+1)}) = \sigma(p^{(j)})$ , and the same is true for the partial quotients. So the same  $k$  applies to  $j + 1$ .

Case 2.  $\sigma(\text{lc}(p^{(j)})) \neq 0$ . Then  $\sigma(\psi_{i_j}) \neq 0$ . As by hypothesis all  $\text{lpp}(g_i)$  remain stable under specialization,  $i_j$  is the first index for which  $\text{lpp}(\sigma(g_{i_j}))$  divides  $\text{lpp}(\sigma(p^{(j)}))$ , so

$$\begin{aligned} \psi'_{i_j} &= \frac{\text{lm}(p'^{(k)})}{\text{lm}(\sigma(g_{i_j}))} \\ p'^{(k+1)} &= p'^{(k)} - \psi'_{i_j} \sigma(g_{i_j}). \end{aligned}$$

Using the induction hypothesis for  $j$  and specializing equations (4) yields:

$$\begin{aligned}\sigma(\psi_{i_j}) &= \frac{\sigma(\text{lm}(p^{(j)}))}{\sigma(\text{lm}(g_{i_j}))} = \frac{\text{lm}(p^{(k)})}{\text{lm}(\sigma(g_{i_j}))} = \psi'_{i_j} \\ \sigma(p^{(j+1)}) &= p^{(k)} - \psi'_{i_j} \sigma(g_{i_j}) = p^{(k+1)},\end{aligned}$$

so in this case we pick  $k + 1$ . Note that the partial quotients also remain stable.  $\square$

The pseudo-division in  $R(\bar{a})[\bar{x}]$  introduces coefficients with denominators in  $R[\bar{a}]$ . But these denominators arise only from leading coefficients of the divisors  $g_1, \dots, g_s$ , and are divisors of a product of powers of them. Under the hypothesis of theorem 1, we can still multiply  $f$  by a convenient factor  $\mu$  in  $R[\bar{a}]$  to eliminate non-vanishing denominators for the specification<sup>‡</sup>. This will only change the normalization of  $f$ .

Writing  $q'_i = \mu q_i$  and  $r' = \mu r$ , we have

$$\mu f = \sum_{i=1}^s q'_i g_i + r'. \quad (5)$$

The result will be a division with quotients  $q'_i$ , ( $1 \leq i \leq s$ ) and remainder  $r'$  in  $R[\bar{a}][\bar{x}]$ . If the condition on the leading monomials is satisfied, theorem 1 guarantees that the result specializes properly except for the not-null factor  $\sigma(\mu) \neq 0$ . We denote this algorithm PDIV and the remainder

$$r' = \overline{f}^{[g_1, \dots, g_s]}.$$

## 4. Buchberger's algorithm and specialization

We denote GB0 the ordinary Buchberger's algorithm (see Cox, Little & O'Shea (1992)), that only add remainders of  $S$ -polynomials, acting on  $k[\bar{x}]$ , where  $k$  is any field (in particular it can be  $R(\bar{a})$  or  $K$ ).

We define PGB0 as the version of GB0 that uses equation (2) for computing  $S$ -polynomials and the normalization of the divisions described in section 3 to avoid denominators when computing in  $R(\bar{a})[\bar{x}]$ . We have:

**THEOREM 2:** *Let  $F \subset K[\bar{a}][\bar{x}]$ ,  $G_0 = \text{PGB0}(F, \succ_{\bar{x}})$  and*

$$W = \text{FACVAR}(\{\text{lc}(g) : g \in G_0\}, \succ_{\bar{a}}).$$

*Then*

- (i)  $G_0$  is a Gröbner basis of  $\langle F \rangle R(\bar{a})[\bar{x}]$  whose polynomials belong to  $R[\bar{a}][\bar{x}]$ .

<sup>‡</sup>There are several possible definitions of  $\mu$  depending on the ring, field and implementation.

- (ii) If  $\sigma(w) \neq 0$ , for all  $w \in W$ , then  $\sigma(G_0)$  is a Gröbner basis of  $\langle \sigma(F) \rangle K[\bar{x}]$ .
- (iii)  $G_0$  can be minimized and reduced to a new basis  $G$  whose elements are in  $R[\bar{a}][\bar{x}]$ . Furthermore, under the above conditions on  $\sigma$ ,  $\sigma(G)$  is the reduced Gröbner basis of  $\langle \sigma(F) \rangle K[\bar{x}]$  (apart from normalization).

We write down the algorithm PGB outlined in theorem 2 (iii). The procedures MINIMIZE and REDUCE to be used are those described in Cox, Little & O’Shea (1992) (with PDIV as division algorithm). The goal of NORMALIZE is ensuring the uniqueness of the reduced Gröbner basis. It takes the primitive part and chooses in a unique form the  $R$ -factor unity in the leading coefficient. The choose depends on the ring and the implementation.

### PGB (Parametric Gröbner Basis)

Input:  $F = \{f_1, \dots, f_s\} \subset R[\bar{x}, \bar{a}]$ , and  $\succ_{\bar{x}\bar{a}}$   
Output:  $G$  the unique reduced Gröbner basis of  $\langle F \rangle$ , with respect to  $[\bar{x}]$ .  
 $W$ : A singular variety outside of which the basis specializes.

$$G_1 := \text{PGB0}(F, \succ_{\bar{x}})$$

$$W := \text{FACVAR}(\{\text{lc}(g, \succ_{\bar{x}}) : g \in G_1\})$$

$$G := \text{NORMALIZE}(\text{REDUCE}(\text{MINIMIZE}(G_1, \succ_{\bar{x}}), \succ_{\bar{x}}), \succ_{\bar{x}\bar{a}}).$$

Using definition 2 and theorem 1 the proof of the theorem above is easy.

The output of PGB are  $G$  and  $W$ . Nevertheless,  $W$  is a sufficient singular variety but not a minimal one. It is important to obtain a minimal solution. This objective will be approached in the next section and solved in section 10.

## 5. GGE: Generalized Gaussian Elimination algorithm

Using PGB produces the unique reduced Gröbner basis. Instead, the computed singular variety is algorithm depending on the development of PGB0 (see Example 5.1).

To reduce the variety as much as possible we introduce a Generalized Gaussian Elimination algorithm (GGE) to transform the initial basis  $F$  of the ideal into a more convenient basis  $F'$  and using it as a pre-processing for PGB0.

The idea of the algorithm comes from ordinary Gaussian elimination, and produces a triangulation of the leading terms with respect to the order  $\succ_{\bar{x}\bar{a}}$ .

### GGE (Generalized Gaussian Elimination)

Input:  $F = [f_1, \dots, f_s] \subset R[\bar{x}, \bar{a}]$ , ( $\bar{x}$  and  $\bar{a}$  as variables)  
 $\succ_{\bar{x}\bar{a}}$  the product elimination order.  
Output:  $F'$  a new basis of  $\langle F \rangle$  that specializes for any  $\sigma$ .

$$G := F; \quad G' = \phi$$

```

WHILE  $G \neq G'$ 
  FOR  $i$  from 1 to  $\#G$  DO
     $G' := G$ 
     $G := \phi$ 
    FOR  $k$  from 1 to  $\#G'$  DO
      IF  $k \neq i$  THEN
         $f := \overline{g}_k^{[g'_i]}$  respect to the order  $\succ_{\overline{x}\overline{a}}$ .
        IF  $f \neq 0$  THEN  $G := G \cup \{f\}$ 
        ELSE  $G := G \cup \{g'_i\}$ 
   $F' := \text{primpart}(G)$ .

```

PROPOSITION 3: *Let  $F \subset R[\overline{a}][\overline{x}]$ , and  $F' := \text{GGE}(F, \succ_{\overline{x}\overline{a}})$ . Then*

- (i)  $F'$  is a new basis of  $\langle F \rangle \text{Quot}(R)[\overline{x}, \overline{a}]$  that has less or equal number of elements than  $F$ .
- (ii)  $\langle \text{lpp}(F, \succ_{\overline{x}\overline{a}}) \rangle \subseteq \langle \text{lpp}(F', \succ_{\overline{x}\overline{a}}) \rangle$ , and also  $\langle \text{lpp}(F, \succ_{\overline{x}}) \rangle \subseteq \langle \text{lpp}(F', \succ_{\overline{x}}) \rangle$ .
- (iii)  $F'$  specializes, i.e.  $\sigma(F')$  is also a basis of  $\langle \sigma(F) \rangle$ .
- (iv) The algorithm terminates.
- (v) Each  $p \in F'$ , is the remainder of dividing  $p$  by  $F' - \{p\}$  with respect to the given order  $\succ_{\overline{x}\overline{a}}$ . We say that  $F'$  is reduced for this order.

*Proof:* (i) At each division in the algorithm we substitute, in a given  $F$ -basis, two polynomials in  $R[\overline{a}][\overline{x}]$ , say  $\{f, g\}$  by  $\{g, r\}$ , where  $f = gh + r$ . As  $f$  and  $g$  can be expressed as a linear combination of  $g$  and  $r$  with coefficients in  $\text{Quot}(R)$ , then  $\{g, r\}$  is a new basis of  $\langle f, g \rangle \text{Quot}(R)[\overline{x}, \overline{a}]$  and, consequently,  $\{g, r\}$  can replace  $\{f, g\}$  to form a new basis of  $F$ . This is the only repeated action in computing the GGE. Observe also, that the final number of polynomials is less or equal than the number of polynomials in  $F$ , as we replace at each step two polynomials by two or one (when the remainder is 0) polynomials.

(ii) The first part is obvious, as the new polynomials are remainders with respect to the order  $\succ_{\overline{x}\overline{a}}$ . When only the  $\overline{x}$  are taken as variables, the proposition is a consequence of  $\succ_{\overline{x}\overline{a}}$  being a compatible elimination order.

(iii) As  $\sigma$  is an homomorphism, we have  $\sigma(f) = \sigma(g)\sigma(h) + \sigma(r)$  and the argument given in (i) can be translated to  $\{\sigma(f), \sigma(g)\}$  for any specialization. It must be noted that in specializing  $F'$ , any coefficient (even the leading one) can become zero. Nevertheless the specialized basis  $\sigma(F')$  is still a basis of  $\langle \sigma(F) \rangle$ .

(iv) We have to prove that the algorithm terminates. Let  $F_i$  be the base after the  $i^{\text{th}}$  loop. Define

$$d_i = \sum_{f \in F_i} \text{multideg}(f, \succ_{\overline{x}\overline{a}}),$$

where  $\text{multideg}(f, \succ)$  is the list of the exponents of the variables of  $\text{lpp}(f, \succ)$ . As  $\#F_i \leq \#F_{i-1}$  and for every division in the algorithm, whenever the leading

term is divisible, the multideg of the remainder is strictly smaller than that of the dividend: necessarily  $d_i \preceq_{\bar{x}a} d_{i-1}$ . If  $d_i = d_{i-1}$  then no leading monomial has changed in loop  $i$ . So, no monomial of a  $f \in F_i$  is divisible by any leading monomial of an element of  $F_i$  (with constant leading coefficient, when we only consider the  $\bar{x}$  as variables), as all possible divisions have been performed in loop  $i$ . The algorithm stops in this case. As  $\succ_{\bar{x}a}$  is a well-order,  $d_i$  necessarily stabilize, and the algorithm effectively terminates. GGE works in the direction of the reduced Gröbner basis, but may not reach it, as not all  $S$ -polynomials are considered and tested.

(v) As all leading coefficients are constants of  $R$ , all polynomials in  $F'$  have been considered as divisors of the others, and no polynomial will be divisible by the remaining. This property is the triangulation property of GGE.  $\square$

We include GGE as a preprocessing step for PGB algorithm, before calling PGB0, in order to transform the initial basis. The resulting singular variety becomes smaller and, in simple examples it suffices to obtain the minimal singular variety.

GGE algorithm is remarkable for linear systems with parameters. Applying it by dividing only with respect to the  $\bar{x}$  variables and divisors with constant leading coefficients, it reduces to the ordinary Gaussian elimination. But applying it, as defined, dividing with respect to variables and parameters, there is no restriction about the leading coefficients of the divisors, and then it provides a perfect first step for a reduced case discussion, as can be seen in the example below.

This algorithm is used in our general DISPGB algorithm. Nevertheless it is interesting by itself, as it provides a triangulation of the basis that is useful in applications.

### 5.1. Example

Consider the following linear system in the variables  $(x, y, z, u)$  with the parameters  $(a, b)$ :

$$F := [ax + 2y + 3z + u - 6, x + 3y - z + 2u - b, \\ 3x - ay + z - 2, 5x + 4y + 3z + 3u - 9].$$

We take  $F$  to be the basis of an ideal in  $\mathbb{Z}[x, y, z, u, a, b]$  and compute a new basis with GGE algorithm, relative to the order  $\text{lex}(x, y, z, u, a, b)$ . The result is:

$$F' := \text{GGE}(F, \text{lex}(x, y, z, u, a, b)) \\ = [756x - 39ab - 4b - 155 - 117a + (117a + 51)u, \\ 189y + 6ab - 107 - 43b + 18a - (18a - 123)u, \\ 756z - 1439 + 236b + 99a + 33ab - (99a - 15)u, \\ (9a^2 - 30a + 21)u - 9a^2 - 3a^2b + 11ab + 22a - 49 + 28b].$$

As we see, the new basis is completely Gaussian reduced:

1. The new basis is triangular: each leading monomial is different when we consider as variables all  $x, y, z, u, a, b$ . In this case, it is also true if we consider only  $x, y, z, u$  as variables.
2. The three first pivots (leading coefficients) of the variables  $x, y, z$  are constants, and only the last leading coefficient in the variable  $u$  is a polynomial in the parameters. It is essentially the singular variety:  $9a^2 - 30a + 21 = 3(a - 1)(3a - 7)$ , precisely the value of the determinant.
3. Instead, the variable  $u$  is not eliminated in the equations giving  $x, y, z$ .
4. In this form, the system is very simple to discuss and to solve.
5. The result is, in this case, a minimal Gröbner basis, but not the reduced one.
6. Moreover, in this example,  $F'$  is itself a comprehensive Gröbner basis.

Now we compute  $\text{PGB}(F', \text{lex}(x, y, z, u), \text{lex}(a, b))$  to obtain the reduced Gröbner basis and the singular variety. We obtain:

$$\begin{aligned}
G &= (9a^2 - 30a + 21)x - 6ab + 9a - 2b - 1, \\
&\quad (9a^2 - 30a + 21)y + 3ab + 20 - 23b, \\
&\quad (9a^2 - 30a + 21)z + 53a - 5ab - 18a^2 + 3a^2b - 39 + 6b, \\
&\quad (9a^2 - 30a + 21)u - 9a^2 - 3a^2b + 11ab + 22a - 49 + 28b]
\end{aligned}$$

$$W = \{a - 1, 3a - 7\}.$$

In this example,  $W$  determines the minimal singular variety, as it includes just the factors of the determinant of the system, which we know by linear algebra to be the singular condition. The general discussion of all cases can be done with DISPGB described in the next sections. The complete output provided by DISPGB for this example can be found in subsection 11.5

If we compute the Gröbner basis using PGB algorithm without previous reduction of  $F$  to  $F'$  by GGE, the basis  $G$  turns out to be the same, but the variety becomes  $W_1 = \{a, a^2 + 35, 3a - 2, a - 1, 3a - 7\}$ , showing the reduction obtained by GGE.

## 6. Specification of specializations

Following Weispfenning (1992), we consider the following families of specializations (conditions in the paper of reference):

**DEFINITION 4:** *Let  $\sigma$  be a specialization and  $N = \{p_1, \dots, p_s\} \subset R[\bar{a}]$  and  $W = \{q_1, \dots, q_r\} \subset R[\bar{a}]$  be sets of polynomials. We say that  $\sigma \in \Sigma(N, W)$ , or that  $(N, W)$  is the actual specification of  $\sigma$  if*

- (i) *All the  $p_i$ 's specialize to zero:  $\sigma(p_i) = 0$ , for  $p_i \in N$ .*
- (ii) *All the  $q_i$ 's specialize to non-zero:  $\sigma(q_i) \neq 0$ , for  $q_i \in W$ .*

We say that  $N$  are the null conditions and  $W$  the not-null conditions of the specification of all  $\sigma \in \Sigma(N, W)$ .

**PROPOSITION 5:** *Let  $\sigma \in \Sigma(N, W)$  be a specialization. If  $f \in \sqrt{\langle N \rangle}$ , (the radical of the ideal  $\langle N \rangle$ ), then  $\sigma(f) = 0$ . Equivalently:*

$$\sigma \in \Sigma(N, W) \implies \sigma \in \Sigma(\sqrt{\langle N \rangle}, W).$$

*Proof:* Let  $\sigma \in \Sigma(N, W)$  and  $f \in \sqrt{\langle N \rangle}$ . Then for some  $n \in \mathbb{N}$  is  $f^n \in \langle N \rangle$ . Thus  $\sigma(f^n) = (\sigma(f))^n = 0$ , and the result follows.  $\square$

**PROPOSITION 6:** *Let  $K$  be an algebraically closed field and  $\Lambda = \Sigma(N, W)$ . Set  $h = \prod_{q \in W} q$ . Then  $\Lambda \neq \emptyset$  iff  $\langle N \rangle \neq K[\bar{a}]$  and  $h \notin \sqrt{\langle N \rangle}$ .*

*Proof:*  $[\implies]$ : By proposition 5.

$[\impliedby]$ : By Hilbert's Nullstellensatz,  $\mathbb{V}(N) = \sqrt{\langle N \rangle}$ , and if  $h \notin \sqrt{\langle N \rangle}$ , it exists  $a_0 \in \mathbb{V}(N)$  for which  $h(a_0) \neq 0$ . Thus  $\sigma_{a_0} \in \Lambda$ , and  $\Lambda \neq \emptyset$ .  $\square$

The set of values of the parameters described by  $\Sigma(N, W)$  is given by

$$\mathbb{V}(N) - [\mathbb{V}(N) \cap [\cup_{q \in W} \mathbb{V}(q)]] .$$

**PROPOSITION 7:** *Let  $W = \{q_1, \dots, q_r\}$ ,  $N$  be a Gröbner basis relative to the order  $\succ_{\bar{a}}$  and  $W' = \text{FACVAR}(\{\bar{q}_1^N, \dots, \bar{q}_r^N\})$ . Then*

$$\Sigma(N, W) = \Sigma(N, W').$$

*Proof:* We have  $q_i = n_i + \bar{q}_i^N$ , where  $n_i \in \langle N \rangle$  and  $\bar{q}_i^N$  is the remainder of the division of  $q_i$  by the basis  $N$ . As  $\sigma$  is an homomorphism,

$$\sigma(q_i) = \sigma(n_i) + \sigma(\bar{q}_i^N) = \sigma(\bar{q}_i^N).$$

So  $\sigma(q_i) \neq 0$  iff  $\sigma(\bar{q}_i^N) \neq 0$ , and we can finally decompose into irreducible factors.  $\square$

**DEFINITION 8 (QUASI-CANONICAL REPRESENTATION OF A SPECIFICATION):**  
A representation  $(N, W)$  is said to be  $k$ -quasi-canonical, if it satisfies

- (i)  $N$  is the reduced Gröbner basis in  $\text{Quot}(R)[\bar{a}]$  for the order  $\succ_{\bar{a}}$  of the set of polynomials that specialize to zero.
- (ii) The polynomials in  $W$  specializing to non-zero are reduced modulo  $N$  and are irreducible over  $k[\bar{a}]$ , where  $k$  is some intermediate field between  $\text{Quot}(R)$  and the algebraic closure  $\bar{K}$ . They are normalized in a canonical form in order to be recognized when compared. (Normalization depends on the field  $k$ , but is easy to define for  $\mathbb{Q}, \mathbb{R}$  and  $\mathbb{C}$ .)
- (iii)  $\prod_{q \in W} q \notin \sqrt{\langle N \rangle}$ .

- (iv) *The polynomials in  $N$  are square-free.*
- (v) *If some  $p \in N$  factors, then no factor of  $p$  belongs to  $W$ .*

PROPOSITION 9: *Any specification that is non-empty in the algebraic closure has a  $k$ -quasi-canonical representation  $(N, W)$ .*

It seems that imposing  $N$  to be a Gröbner basis of  $\sqrt{\langle N \rangle}$ , can provide a unique representation. Nevertheless, for practical computation reasons, we do not impose  $N$  to be radical.

*Proof:* (i) We can choose a Gröbner basis to define the ideal  $\langle N \rangle$ . This will report benefits in the algorithms.

- (ii) By proposition 7 we can reduce the polynomials in  $W$  modulo  $N$ , decompose into irreducible factors and normalize them (depending on the field).
- (iii) By proposition 6,  $\prod_{q \in W} q \notin \sqrt{\langle N \rangle}$  is a necessary and sufficient condition for  $N$  and  $W$  to be compatible conditions in the algebraic closure, i.e. to determine a non-empty  $\Sigma(N, W)$ .
- (iv) We can drop any multiple factor in  $N$  as a consequence of proposition 5.
- (v) We can also drop any factor of a polynomial  $p \in N$  belonging to  $W$  as  $p = gh$  and  $g \in W$  imply  $\sigma(p) = 0$  and  $\sigma(g) \neq 0$ , which leads to  $\sigma(h) = 0$ .

□

If in the development of the algorithms a polynomial  $p' \in R[\bar{a}]$  such that  $p' \in \sqrt{\langle N \rangle}$  and  $p' \notin \langle N \rangle$  is detected, we can refine the representation of the specification adding  $p'$  to the base  $N$ , and re-computing the new Gröbner basis  $N'$ . We then have:

$$\langle N \rangle \subseteq \langle N' \rangle \subseteq \sqrt{\langle N \rangle}.$$

Then  $(N', W)$ ,  $(N, W)$  and  $(\sqrt{\langle N \rangle}, W)$  are representations of the same specification  $\Sigma(\sqrt{\langle N \rangle}, W)$ , but  $(N', W)$  is a finer representation than  $(N, W)$ . Even if some  $p \in N'$  has a factor in  $W$ , it can also be dropped, as it specializes to non-zero.

DEFINITION 10: *Let  $f_1, f_2 \in R[\bar{a}][\bar{x}]$ . We say that  $f_1 \approx_\sigma f_2$  (are  $\sigma$ -equivalents) iff  $\sigma(f_1) = \sigma(f_2)$ .*

When a specification is assumed, then we can substitute the polynomials in a set by  $\sigma$ -equivalent polynomials. This will be done all along the algorithm. It must be noticed, that doing so, the new  $\sigma$ -equivalent set is no more a basis of the original  $F \subset R[\bar{a}][\bar{x}]$ . This is an important observation for the understanding of the algorithm. Doing so, we abandon to compute a CGB, but the algorithm is simplified towards our objective. Observe that by doing this, unnecessary polynomials specializing to zero in the actual specification are also dropped.

DEFINITION 11: *Given two specifications  $(N_1, W_1)$  and  $(N_2, W_2)$  we say that  $(N_2, W_2) \geq (N_1, W_1)$ , if  $\sqrt{\langle N_1 \rangle} \subseteq \sqrt{\langle N_2 \rangle}$  and  $W_1 \subseteq W_2$ .*

*If  $(N_2, W_2) \geq (N_1, W_1)$  and  $\sqrt{\langle N_1 \rangle} \neq \sqrt{\langle N_2 \rangle}$  or  $W_1 \neq W_2$ , then we say that  $(N_2, W_2) > (N_1, W_1)$ .*

DEFINITION 12: *The specifications where  $N = \{0\}$  are denoted generic, in the sense that  $\Sigma(\{0\}, W)$  contains all specializations except a set in the  $m - 1$  dimensional variety  $\cup_{q \in W} \mathbb{V}(q)$  defined by  $W$ .*

### 6.1. CANSPEC algorithm

The algorithms will only use quasi-canonical representations of specifications.

PROPOSITION 13: *Given the specification  $(N, W)$ , the algorithm CANSPEC produces a  $k$ -quasi-canonical representation  $(N', W')$  if  $\text{test} = \text{true}$ . If  $\text{test} = \text{false}$  then  $(N, W)$  are incompatible, i.e.  $\Sigma(N, W) = \phi$*

#### CANSPEC

Input:  $(N, W)$ : the specification of  $\sigma$ .  
 $\succ_{\bar{a}}$ : monomial order for the parameters  
Output:  $\text{test}$ : has the value *true* if  $\Sigma(N, W) \neq \phi$  in the algebraic closure and *false* if they are incompatible.  
 $(N', W')$ : A  $k$ -quasi-canonical representation of the specification.

```

 $W' := \text{FACVAR}(\{\bar{q}^N : q \in W\})$ 
 $\text{test} := \text{true}; t := \text{true}; N' := N$ 
 $h := \prod_{q \in W} q$ 
IF  $h \in \sqrt{\langle N' \rangle}$  THEN  $\text{test} := \text{false}; N' := \{1\}$ ; STOP
WHILE  $t$  DO
   $t := \text{false}$ 
   $N'' :=$  Drop any factor of a polynomial in  $N'$  that belongs
    to  $W'$ , as well as multiple factors.
  IF  $N'' \neq N'$  THEN
     $t := \text{true}$ 
     $N' := \text{GB}(N'', \succ_{\bar{a}})$ 
     $W' := \text{FACVAR}(\{\bar{q}^N : q \in W'\})$ .

```

*Proof:* The first step reduces  $W'$  modulo  $N$  using proposition 7. Then one uses proposition 6 to test compatibility. It is sufficient to test it only once as no transformation to a new representation  $(N', W')$  will alter it.

Then the loop intends to simplify  $N'$  eliminating any factor in  $N'$  belonging to  $W'$ . If one factor is found then one recomputes the Gröbner basis for  $N'$

and reduces  $W'$  modulo  $N'$ . The loop is finite by the ascending chain condition (ACC).

Thus all the conditions for  $(N', W')$  to be a quasi-canonical representation are met.  $\square$

## 7. DISPGB discussion algorithm

We are now ready to give the general description of the main algorithm DISPGB (Discussing Parametric Gröbner Bases), which is very similar to Weispfenning's construction of a Gröbner system. Its goal is to build up a binary tree structure beginning at the root. Let us give the algorithm:

### DISPGB

Input:  $F \subset R[\bar{x}][\bar{a}]$ , and the orders  $\succ_{\bar{x}}$  and  $\succ_{\bar{a}}$ .

Output: The table  $T[v]$  having tree structure.

$B := \text{GGE}(F, \succ_{\bar{x}\bar{a}})$

$T[\phi] := (\phi, B, \phi, \phi)$  (Creates and stores the root vertex)

BRANCH( $\phi, B, \phi, \phi$ ) (Begins the recursive building of the tree).

At each vertex  $v$  the following objects are stored in the table variable

$$T[v] = (c_v, B_v, N_v, W_v)$$

1. The new condition  $c_v$  of the form  $p(\bar{a}) = 0$  (for type 0 vertices), or  $p(\bar{a}) \neq 0$  (for type 1 vertices), that is assumed when the algorithm gains access to vertex  $v$  in order to increase the specification of  $\sigma$ .
2. The pair  $(N_v, W_v)$  determining a quasi-canonical representation of the specification of  $\sigma$  at this vertex (see definition 8).
3.  $B_v \subset R[\bar{a}][\bar{x}]$ , a set of polynomials that specializes to a basis of the ideal  $\langle \sigma(F) \rangle$  for any  $\sigma \in \Sigma(N_v, W_v)$ .

DISPGB begins by computing  $\text{GGE}(F)$ . As discussed in section 5 this reduces the number of cases to be discussed, reducing at the same time the singular variety.

At the root, the initial basis transformed by GGE is stored, and the rest of arguments are empty. So  $T[\phi] := (\phi, B, \phi, \phi)$ .

Then the recursive algorithm BRANCH, described in section 9, is called: bifurcation takes place making the dichotomic decision about some leading coefficient of a polynomial in the actual basis to become zero or not-zero in the specialization, like in dynamical evaluation (Duval, 1995). The recursive algorithm BRANCH, combined with NEWVERTEX, described also in section 9, will be responsible of the control flow of the procedure. As the algorithm progresses in depth, the the specification of the specialization is refined at the branches,



about the specialization of the leading coefficient to zero or not. The goal of CONDPGB is, given the specification and  $B \subset R[\bar{a}][\bar{x}]$ , where  $\sigma(B)$  is a basis of  $\langle \sigma(F) \rangle$ , to advance in the direction of a specializing Gröbner basis as much as possible.

### 8.1. NEWCOND algorithm

PROPOSITION 14: *Let  $(N, W)$  be the specification of  $\sigma$ , and  $f \in R[\bar{a}][\bar{x}]$ . Then the following algorithm NEWCOND determines the quadruplet  $(cd, f', N', W')$  where  $f' \approx_\sigma f$ .*

*If  $cd = \phi$ , then  $\sigma(\text{lc}(f', \succ_{\bar{x}})) \neq 0$  for  $\sigma \in \Sigma(N, W)$ . If  $cd \neq \phi$ , set  $c = \prod_{w \in cd} w$ . Then  $\sigma(\text{lc}(f', \succ_{\bar{x}}))$  vanishes or not depending on the nullity or not of  $\sigma(c)$ .*

*$(N', W')$  is a refinement of  $(N, W)$ , when some polynomial in  $\sqrt{\langle N \rangle}$  is detected.*

#### NEWCOND

Input:  $f$ :  $\in R[\bar{a}][\bar{x}]$   
 $(N, W)$ : Null and not-null conditions of the specification of  $\sigma$ .  
Output:  $cd$ : The new conditional set.  
 $f'$ : The  $\sigma$ -equivalent to  $f$  with deciding leading coefficient.  
 $(N', W')$ : A specialization refinement of  $(N, W)$ .

```

 $f' := f$ ;
 $test := \text{true}$ 
 $N' := N$ 
WHILE  $test$  DO
  IF  $\text{lc}(f') \in \sqrt{\langle N' \rangle}$  THEN
     $f' := f' - \text{lm}(f')$ 
     $N' := \text{GB}(N' \cup \text{lc}(f'), \succ_{\bar{a}})$ 
     $W' := \{\bar{w}^{N'} : w \in W\}$ 
  ELSE  $test := \text{false}$ 
 $f' := \overline{f'}^{N'}$ 
 $cd := \text{FACVAR}(\text{lc}(f')) - W'$ 

```

*Proof:* If the specification of  $\sigma$  implies that  $\sigma(\text{lc}(f)) = 0$ , NEWCOND begins eliminating iteratively the leading term of  $f$ . In this process, some polynomial in  $\sqrt{\langle N \rangle}$  can be detected. This is used to improve  $N$  and a new Gröbner basis  $N'$  is recomputed. This is then used to improve also  $W$ . When no more leading monomials of  $f'$  can be dropped, then  $f'$  is reduced with respect to  $N'$  dividing by it.  $(N', W')$  is now not sufficient to decide if the actual  $\text{lc}(f')$  specialize to zero as a consequence of the null condition  $N'$ .

The algorithm now computes  $cd := \text{FACVAR}(\text{lc}(f')) - W'$ . Obviously, if  $cd =$

$\phi$  then  $\sigma(\text{lc}(f')) \neq 0$  for  $\sigma \in \Sigma(N', W')$ . If not,  $\sigma(\text{lc}(f', \succ_{\bar{x}}))$  vanishes or not depending on the nullity or not of  $\sigma(c)$ .  $\square$

When NEWCOND has concluded and has detected a new condition, in order to continue the specification of  $\sigma$ , DISPGB will have to decide either

- (0) The product of all polynomials in  $cd$  become zero by specialization ( $c = 0$ ).
- (1) All polynomials in  $cd$  become different from zero by specialization ( $c \neq 0$ ).

## 8.2. CONDPGB: Conditional Parametric Gröbner Basis algorithm

At a given point DISPGB needs to use a Buchberger's type algorithm that takes into account the specification and intends to determine a specializing Gröbner basis. We denote it CONDPGB (Conditional Parametric Gröbner Basis). <sup>¶</sup>

### CONDPGB

Input:  $B$ : The actual specializing basis.  
 $(N, W)$ : The actual specification of  $\sigma$ .  
Output:  $test$ : if  $test = true$  then  $\sigma(B')$  is yet the Gröbner basis.  
 $B'$ : The new completed specializing basis.  
 $(N', W')$ : The specification  $(N, W)$  can be refined in the process.

```

test := true; s := #B; J := {(i, j) : 1 ≤ i < j ≤ s}
B' := B; N' := N; W' := W; t := s
WHILE J ≠ ∅ and test DO
  Select (i, j) ∈ J
  J := J - {(i, j)}
  IF Buchberger's reductibility criterions are false THEN
    S := SPOL(B'[i], B'[j], ≻̄x)B'
    S := S̄x̄aN'
    IF S ≠ 0 THEN
      (cdec, S, N', W') := NEWCOND(S, N', W')
      IF cdec = ∅ THEN
        IF S ≠ 0 THEN
          t := t + 1
          B' := B' ∪ {S}
          J := J ∪ {(i, t) : 1 ≤ i < t}
        ELSE
          test := false
          B' := B' ∪ {S}

```

<sup>¶</sup>Actually, a new release of the implementation is in course of development. It introduces improvements in the algorithm CONDPGB to accelerate Buchberger's algorithm, that are not yet described here.

IF *test* THEN

$$B' := \text{REDGB}(\text{MINGB}(B', \succ_{\bar{x}}, \succ_{\bar{x}}, \succ_{\bar{a}}).$$

**PROPOSITION 15:** *Let  $(N, W)$  be the specification of  $\sigma$  and let  $B \subset R[\bar{a}][\bar{x}]$  be a set of polynomials for which:*

- (i)  $\sigma(B)$  is a basis of  $\langle \sigma(F) \rangle$ .
- (ii)  $\sigma(\text{lc}(g, \succ_{\bar{x}})) \neq 0$  for  $g \in B$  and  $\sigma \in \Sigma(N, W)$ .

The algorithm **CONDPGB** determines the quadruplet  $(\text{test}, B', N', W')$ .

If  $\text{test} = \text{true}$ , then  $\sigma(B')$  is (except for normalization), the reduced Gröbner basis of  $\langle \sigma(F) \rangle$  for  $\sigma \in \Sigma(N', W')$ .

If  $\text{test} = \text{false}$ , then  $B'$  is an extended set of  $B$  for which  $\langle \text{lpp}(B, \succ_{\bar{x}}) \rangle \subsetneq \langle \text{lpp}(B', \succ_{\bar{x}}) \rangle$ , and  $B'$  contains at least one polynomial for which the actual specification  $(N', W')$  cannot decide if its leading coefficient specializes to zero or not.

*Proof:* **CONDPGB** is essentially Buchberger's algorithm (PGB0) with the following differences:

- (i) Step  $S := \overline{S}_{\succ_{\bar{x}\bar{a}}}^{N'}$ .

After computing the usual remainder of the  $S$ -polynomial, the result is reduced with respect to  $N'$  for the order  $\succ_{\bar{x}\bar{a}}$ . Obviously, both polynomials are  $\sigma$ -equivalents, as the difference is a polynomial with coefficients in  $\langle N' \rangle$  that specializes to 0.

- (ii) Step  $(c_{dec}, S, N', W') := \text{NEWCOND}(S, N', W')$ .

If  $S \neq 0$  then, before continuing PGB0, **CONDPGB** applies **NEWCOND** to test if  $\text{lc}(S, \succ_{\bar{x}})$  specializes to zero or not. This can reduce  $S$  to a  $\sigma$ -equivalent polynomial, say  $S'$ , that can eventually be 0. Three cases can occur:

- a) If  $S' = 0$  then there is nothing new to do and the algorithm continues with PGB0.
- b) If  $S' \neq 0$  and  $\sigma(\text{lc}(S, \bar{x})) \neq 0$  then  $S'$  is adjoined to the base and PGB0 continues.
- c) If  $S' \neq 0$  and  $c_{dec} \neq \phi$  then  $S'$  is adjoined to the base and **CONDPGB** stops, returning the new base  $B'$ . It is important to note that, in that case, at least one new polynomial has been adjoined to the base, whose leading power product cannot be divided by the others in  $B'$ . Thus the ideal of leading power products is strictly greater than before.

If **CONDPGB** never goes through the sub-case c), then it continues until a Gröbner basis with non-zero specializing leading coefficients is reached. In this case, **CONDPGB** determines first the minimal and then the reduced Gröbner base. Then  $c_{dec} = \phi$  is returned.

## 9. Control Algorithms □

We give now the algorithms BRANCH and NEWVERTEX that are recursive procedures calling one another. The control flow of the main DISPGB algorithm is governed by them.

### BRANCH

Input:  $v$ : Label of the vertex.  
 $B$ : Specializing basis at the vertex  $v$ .  
 $(N, W)$ : Specification of  $\sigma$  at vertex  $v$  (not necessarily canonical).  
Output: Recursive algorithm. It stores the refined  $(B', N', W')$  (basis and quasi-canonical representation) at the vertex  $v$ , creating two new hanging vertices when necessary or marking the vertex as terminal. It manages the control flow.

```

 $cd := \phi$ 
FOR  $i$  TO  $\#B$  WHILE  $cd = \phi$  DO
   $f := B[i]$ 
   $(cd, f', N', W') := \text{NEWCOND}(f, N, W)$ 
  Substitute the  $i$ -th element  $f$  of  $B'$  by  $f'$ 
 $pivot := i - 1$ 
 $T[v] := (-, B, N', W')$  ( $cond$  is already stored in  $T(v)$ . Refinement of data)
IF  $cd = \phi$  THEN
   $(test, B', N', W') := \text{CONDPGB}(B, N', W')$ 
  IF  $test$  THEN
     $T[v] = (-, B', N', W', \text{terminal vertex})$ 
    STOP ( $B'$  is already the Gröbner basis)
  ELSE
    BRANCH( $v, B', N', W'$ ) (Further refinement is needed)
ELSE (A pair of new hanging vertices is created)
  NEWVERTEX( $1, v, cd, B', N', W', pivot$ )
  NEWVERTEX( $0, v, cd, B', N', W', pivot$ ).

```

BRANCH begins by testing the polynomials in  $B$  using NEWCOND. This can refine the data at the vertex.

If all have leading coefficients specializing to non-zero ( $cd = \phi$ ), then CONDPGB is called in order to find the specializing Gröbner basis. If this objective is reached, the results are stored at the vertex and the procedure stops. Else, at least one new polynomial with non-decided leading coefficient has been adjoined by CONDPGB (see proposition 15). Thus BRANCH is recursively called in order to increase the refinement and advance in the Gröbner basis direction.

Otherwise some polynomial in  $B$  with non-decided leading coefficient is detected ( $cd \neq 0$ ). Then it is taken as pivot and is used to bifurcate the tree (NEWVERTEX) with two new branches of type 0 and 1, increasing the specification.

**NEWVERTEX**

Input:  $n$  0 or 1. If  $n = 1$ , the new condition is taken not-null. If  $n = 0$  then it is taken null.  
 $u$ : Label of previous vertex. Current vertex becomes  $v := (u, n)$ .  
 $cd$ : Irreducible factors of the new condition.  
 $B$ : The current base at previous vertex.  
 $(N, W)$ : Specification of  $\sigma$  at previous vertex.  
 $pivot$ : The index of pivot polynomial of  $B$  whose leading coefficient is responsible of the new condition.

Output: NEWVERTEX creates the new vertex with label  $v$  and stores the new values of  $(cond_v, B_v, N_v, W_v)$  in  $T(v)$ . Then it call BRANCH to continue the process.

```

 $v := (u, n)$ 
 $c := \prod cd$ 
IF  $n = 0$  THEN
   $cond := (c = 0)$ 
   $W' := W$ 
   $N' := \text{GB}(cd \cup N, \succ_{\bar{x}})$ 
ELSE ( $n = 1$ )
   $cond := (c \neq 0)$ 
   $W' := W \cup cd$ 
   $N' := N$ 
   $B' :=$  Substitute every polynomial  $g$  in  $B$  whose  $\text{lpp}(g, \succ_{\bar{x}})$  can be
    divided by  $\text{lpp}(g_{pivot}, \succ_{\bar{x}})$ , by the  $S$ -polynomial of both.
 $(test, N', W') := \text{CANSPEC}(N', W')$ 
IF  $test$  THEN (Specification is compatible)
   $B' := \text{GGE}(\{\bar{g}^{N'} : g \in B'\} - \{0\}, \succ_{\bar{x}})$  (Optional ||)
   $T[v] := (cond, B', N', W')$  (Create vertex and store)
  BRANCH( $v, B', N', W'$ ) (Begin the analysis)
ELSE STOP (Incompatible specification has been detected).
```

**THEOREM 16:** *Given  $F \subset R[\bar{a}][\bar{x}]$  and the monomial orders  $\succ_{\bar{x}}, \succ_{\bar{a}}$ , DISPGB*

<sup>||</sup>The use of GGE at this point of the algorithm is optional. Nevertheless we observed experimentally that in some examples (like in 11.3) it can drastically reduce the size of the tree, intermediate computations and the number of cases. See the footnote of subsection 8.2.

constructs a table  $T$  with binary tree structure that contains at each terminal vertex the quadruple

$$T_v = (cd, B, N, W)$$

where, either the vertex is marked as incompatible, or

- (i)  $(N, W)$  is a quasi-canonical specification of a specialization.
- (ii) For  $\sigma \in \Sigma(N, W)$ ,  $\sigma(B)$  is the reduced Gröbner basis of  $\sigma(F)$  (except for normalization).
- (iii) Any specialization corresponds to one and only one of the specifications of the terminal vertices.
- (iv) The algorithm terminates.

Moreover, traversing the tree from the root and considering the condition  $cd$  at the vertices, a dichotomic discussion of the cases is provided.

*Proof:* (i) Whenever an extended specialization is computed (in NEWVERTEX), the algorithm CANSPEC is called, producing either a quasi-canonical representation of the specification, or detecting incompatible conditions. In that case it marks the vertex as terminal and with incompatible conditions. Thus, only quasi-canonical representations of specializations are used.

(ii) A vertex is marked as terminal only if conditions are incompatible or if CONDPGB has concluded with a basis specializing to the reduced Gröbner basis of  $\sigma(F)$  (except for normalization).

(iii) Is an immediate consequence of the dichotomic character of the decisions taken at each vertex ( $\sigma(c) = 0$  or  $\sigma(c) \neq 0$ ).

(iv) Is a consequence of the ascending chain condition (ACC) of the ideals in Noetherian rings. When BRANCH is called, the leading coefficients in  $B$  are analyzed. If all these leading coefficients specialize to non-zero in the specification, then CONDPGB is called. Proposition 15 ensures that  $\langle \text{lpp}(B, \succ_{ox}) \rangle$  strictly increases in the call.

If the specialization to zero or not of some leading coefficient cannot be decided, then bifurcation takes place calling NEWVERTEX twice. The branch for which the null-condition is assumed, increases the ideal  $\langle N \rangle$ . The branch for which the not-null condition is assumed increases the number of assumed not-null leading coefficients under specialization, without modifying the base. If in the next decisions none of the above cases occur, then it will arrive a moment where all leading coefficients will be assumed not-null in the specification, and then, CONDPGB will be called, increasing  $\langle \text{lpp}(B, \succ_{ox}) \rangle$ .

So in any case we have ascending chains of ideals that stabilize. Consequently the algorithm terminates.  $\square$

## 10. Minimal Singular Variety: GENCASE algorithm.

In section 4, we proved that PGB determines the unique reduced Gröbner basis  $G$  and a singular variety  $W$  such that, outside  $W$ ,  $G$  is specialization-invariant.

This is the only  $m$ -dimensional case. All other cases are contained in the  $(m-1)$ -dimensional variety  $W$ .

Let us describe a case  $C$  obtained by DISPGB by the 5-tuple  $C = (l, p, B, N, W)$ , where  $l$  is the label,  $p$  is the the list of inequalities and equalities in the order that they have been taken (i.e. the path of the case),  $B$  is the reduced Gröbner basis, and  $(N, W)$  the quasi-canonical representation of the specification. Let  $L$  be the list of cases obtained by DISPGB and in the form described above.

Except for the trivial problems where the reduced Gröbner basis specializes everywhere, DISPGB always obtains the case  $C_0$ , labeled  $[1, \dots, 1]$ , specified by  $(N_0 = \phi, W_0)$ , corresponding to the PGB solution. The inequalities  $W_0$  are sufficient to ensure proper specialization of  $B_0$ . But not all the conditions in  $W_0$  are always necessary: some other cases can also specialize properly.

A case  $C = (l, p, B, N, W)$  in  $L$  is said to be *special* if  $\text{lpp}(B) \neq \text{lpp}(B_0)$  and *normal* if  $\text{lpp}(B) = \text{lpp}(B_0)$ . Let  $L_s$  denote the set of special cases in  $L$  and  $L_n$  the set of normal cases in  $L$  (including  $C_0$ ). Using these notations we can define the minimal variety and state a theorem about it.

**DEFINITION 17:** *Given a set of polynomials  $F \subset R[\bar{a}][\bar{x}]$ , the minimal singular variety  $V_{\min}$  is the minimal variety of  $K^m$  containing all special cases  $L_s$  provided by DISPGB.*

**THEOREM 18:** *Given  $F \subset R[\bar{a}][\bar{x}]$ , let  $L$  be the list of cases determined by DISPGB,  $C_0$  the case corresponding to the label  $[1, \dots, 1]$ ,  $L_n \subseteq L$  the set of normal cases,  $L_s \subseteq L$  the set of special cases and*

$$W_{0m} = \text{FACVAR}(\text{lc}(B_0, \succ_{\bar{x}})).$$

*Then*

- (i) *The minimal singular variety  $V_{\min} = \bigcup_{w \in W_{\min}} \mathbb{V}(w)$ , determined by its irreducible components, verifies:  $W_{0m} \subseteq W_{\min} \subseteq W_0$ .*
- (ii) *For all normal cases, the Gröbner basis is specialization-invariant.*

*Proof:* (i) Conditions  $W_0$  are sufficient for properly specialization of the algorithms. But some of them may be unnecessary. On the other hand, the conditions in  $W_{0m}$  are obviously necessary in order to have a normal case. Thus the components of  $V_{\min}$  are those of  $W_{0m}$  plus, at most, some of the irreducible components in  $W_0$  not in  $W_{0m}$ .

- (ii) Let  $G$  be the reduced Gröbner basis of the ideal  $I = \langle F \rangle R(\bar{a})[\bar{x}]$ . Under the hypothesis  $W_0$ , we have  $\sigma_0(G) = B_0$  (except for normalization) as all steps of the computation of  $G$  specialize properly to  $B_0$ . So

$$\langle \text{lpp}(I) \rangle = \langle \text{lpp}(G) \rangle = \langle \text{lpp}(B_0) \rangle = \langle \text{lpp}(\langle \sigma_0(F) \rangle) \rangle.$$

But by the definition,  $\langle \text{lpp}(B_{L_{ni}}) \rangle = \langle \text{lpp}(B_0) \rangle$  for all cases in  $L_n$ . As

$B_{L_{ni}}$  is the reduced Gröbner basis for the case  $L_{ni}$ , we have  $\langle \text{lpp}(B_{L_{ni}}) \rangle = \langle \text{lpp}(\langle \sigma_{L_{ni}}(F) \rangle) \rangle$ . Thus  $\langle \text{lpp}(I) \rangle = \langle \text{lpp}(\langle \sigma_{L_{ni}}(F) \rangle) \rangle$ , so that, for all cases in  $L_n$ , proper specialization holds.

□

Using part (ii) of theorem 18 we define the *generic case*  $C_g$  by the specification  $(N_g, W_g) = (\phi, W_{\min})$ . It contains all the normal cases  $L_n$  that are completely outside  $V_{\min}$ . There can exist cases in  $L_n$  that have a non-empty intersection with  $V_{\min}$ . The parts of these cases outside  $V_{\min}$  are in the generic case, and the parts inside  $V_{\min}$  are considered as special cases, even if they specialize properly.

This theorem does not determine completely  $W_g$ , although it limits quite strictly the candidates. In practice, we observe that in most examples  $W_{\min} = W_{0m}$ , as was pointed out by V. Weispfenning, in the discussions. Nevertheless there are examples where  $W_{\min}$  is strictly greater than  $W_{0m}$ .\*\*

This induces GENCASE algorithm to improve the output of DISPGB.

### GENCASE

Input:  $T$  The table constructed by DISPGB.  
 $\succ_{\bar{x}}$ :  $\bar{x}$ -termorder.  
 $\succ_{\bar{a}}$ :  $\bar{a}$ -termorder  
Output:  $L'$  The list of cases containing:  
the generic case  $C_g = (\text{'Generic case'}, p_g, B_g, N_g = \phi, W_g)$ ,  
and the normal and special cases inside  $W_g$ .

```

L := {C(l) : l ∈ terminal vertices of T }
C0 := Select from L the case labeled[1..1] specified by (N0 = φ, W0)
L1 := L - C0
lp0 := {lpp(f) : f ∈ B0}
L0 := Select from L1 the cases having lp0 as set of lpp(f)
Ls := L1 - L0
Wg := FACVAR(lc(B0,  $\succ_{\bar{x}}$ ))
Vg =  $\cup_{w \in W_g} \mathbb{V}(w)$ 
WHILE not all cases in Ls belong to Vg DO
  Warning!
  Add convenient elements of W0 - Wg to Vg
Cg := ( 'Generic case' , {w ≠ 0 : w ∈ Wg} , B0, φ, Wg )
Lns := Select all cases in L0 with non-empty intersection with Vg and
intersect them with Vg
L' := Cg ∪ Lns ∪ Ls.

```

GENCASE begins testing if all special cases obtained by DISPGB are inside

\*\*At the end of the tutorial joint to the implemented software, such an example is presented.

$W_{0m}$ . If so, the generic case substitutes all cases in  $L_n$ . Those being completely outside  $V_{\min}$  will disappear from the list, and those having a non-empty intersection with  $V_{\min}$  will be restricted to their intersection with  $V_{\min}$  and, consequently, this condition will be added to their specification. Those cases will be considered special. If not, GENCASE gives a warning and adds convenient polynomials of  $W_0 - W_{0m}$  to  $V_{\min}$  to reach the goal.

For linear systems the system determinant gives the generic case condition. The corresponding generalized value for general polynomial systems is thus

$$\Delta = \prod_{w \in W_g} w.$$

It must be pointed out that the given definition of minimal singular variety is algorithm depending. Although this concept seems to be an intrinsic object, no proof of its algorithm independence is provided here. This remains an open question.

## 11. Applications

Many interesting applications can be examined using DISPGB and GENCASE with very satisfactory results.

### 11.1. Linear systems

A particularity of DISPGB is its exceptional behaviour for linear systems. Let us give an example. Consider the following linear system with the variables  $(x, y, z)$  and the parameters  $(a, b, c)$ :

$$\begin{aligned} x + cy + bz + a &= 0 \\ cx + y + az + b &= 0 \\ bx + ay + z + c &= 0 \end{aligned}$$

Applying DISPGB and GENCASE with respect to the monomial orders  $\text{lex}(x, y, z)$  and  $\text{lex}(a, b, c)$ , the following polynomial appears in the discussion:

$$\Delta \equiv a^2 + b^2 + c^2 - 2abc - 1,$$

that is nothing else than the value of the determinant of the system. Table 1 resumes the discussion of the cases.

GENCASE summarizes the  $[1,1]$  case and the  $[0,1]$  cases obtained by DISPGB into the generic case  $\Delta \neq 0$ , as it corresponds to the discussion by determinants. The  $[1,1]$  case corresponds to a larger singular variety, namely  $\mathbb{V}(\Delta) \cup \mathbb{V}(c^2 - 1)$ , and the  $[0,1]$  case corresponds to  $c^2 - 1 = 0, a - bc \neq 0$ . But the final use of GENCASE, reduces both cases to only one, and the singular variety to the minimal one. All other are special cases corresponding to  $\Delta = 0$ .

**Table 1:** Discussion of the linear system. (Section 11.1).

Case	Conditions	Gröbner basis
Generic case	$\Delta \neq 0$	$[\Delta z + c^3 - b^2c - a^2c + 2ab - c,$ $\Delta y + b^3 - c^2b - a^2b + 2ca - b,$ $\Delta x + a^3 - c^2a - b^2a + 2bc - a]$
Special cases	$\Delta = 0$	
[1,0,1]	$c^2 - 1 \neq 0,$ $\Delta = 0,$ $ab - c \neq 0$	[ 1 ]
[1,0,0]	$c^2 - 1 \neq 0,$ $\Delta = 0,$ $ab - c = 0$	$[(c^2 - 1)y + (bc - a)z - b^3 + c^2b,$ $(c^2 - 1)x + (ca - b)z + cb^3 - a + c^2a - c^3b]$
[0,0,1]	$c^2 - 1 = 0,$ $a - bc = 0,$ $b^2 - 1 \neq 0$	[ $z + c,$ $x + cy$ ]
[0,0,0]	$c^2 - 1 = 0,$ $a - bc = 0,$ $b^2 - 1 = 0$	[ $x + cy + bz + bc$ ]

One can see that the discussion provided by the algorithm cannot be improved: only strictly different cases corresponding to different sets of leading power products of the reduced Gröbner basis appear. It is hard to obtain such a reduced discussion using only determinants.

### 11.2. Singular points of a conic

We study the singular points of the conic

$$x^2 + by^2 + 2cxy + 2dx + 2ey + f = 0.$$

The polynomial system is:

$$[x^2 + by^2 + 2cxy + 2dx + 2ey + f, x + cy + d, by + cx + e].$$

The following “discriminant” appears in the discussion

$$\Delta = bd^2 - bf + c^2f - 2ecd + e^2.$$


Table 2 describes the discussion. The generic case corresponds to non-degenerate conics. Case [0,1] corresponds to two parallel lines. This is an interesting case, as even if the Gröbner basis specializes properly, it is inside the singular variety, corresponding to a special case (degenerate conic without singular points). Finally, cases [1,0] and [0,0,1] correspond to two incident lines, and case [0,0,0] to two coincident lines.

### 11.3. The inverse kinematics problem for a simple robot

We consider now the simple plane robot of fig. 2 with two arms of lengths 1 and  $l$  respectively.

**Table 2:** Discussion of singular points of a conic. (Section 11.2).

Case	Conditions	Gröbner basis
Generic case	$\Delta \neq 0$	[ 1 ]
Special cases	$\Delta = 0$	
[0, 1]	$cd - e = 0, d^2 - f \neq 0$	[ 1 ]
[1, 0]	$cd - e \neq 0$	[ $d^2 - f + (cd - e)y, -de + cf + (cd - e)x$ ]
[0, 0, 1]	$cd - e = 0, d^2 - f = 0,$ $b - c^2 \neq 0$	[ $y, x + d$ ]
[0, 0, 0]	$cd - e = 0, d^2 - f = 0,$ $b - c^2 = 0$	[ $x + cy + d$ ]

**Figure 2:** Simple two arms robot.

The inverse kinematics problem is provided by the solution of the following system:

$$\begin{aligned}
 r - c_1 + l(s_1 s_2 - c_1 c_2) &= 0 \\
 z - s_1 - l(s_1 c_2 + s_2 c_1) &= 0 \\
 s_1^2 + c_1^2 - 1 &= 0 \\
 s_2^2 + c_2^2 - 1 &= 0.
 \end{aligned}$$

where  $(s_1, c_1, s_2, c_2)$  are the sines and cosines of the two angles  $\theta_1$  and  $\theta_2$  defined in fig. 2. In the inverse kinematics problem these are the variables to be determined for the parameter values  $(r, z, l)$ . Applying DISPGB plus GENCASE to this system, using  $\text{lex}(s_1, c_1, s_2, c_2)$  and  $\text{lex}(r, z, l)$  orders, the discussion provided in table 3 is obtained.

DISPGB provides  $V = \mathbb{V}(l) \cup \mathbb{V}(r) \cup \mathbb{V}(r^2 + z^2) \cup \mathbb{V}(z) \cup \mathbb{V}(-l^2 + r^2 + 1)$  as singular variety, plus 4 other cases that are in fact special cases of the generic case. GENCASE obtains the minimal singular variety  $V_g = \mathbb{V}(l) \cup \mathbb{V}(r^2 + z^2)$  in only one generic case containing all previous five cases.

Moreover, the minimal singular variety has a simple geometrical interpretation: the special cases correspond either to the end of the robot being at the origin, or to the length of the second arm being 0.

The special cases [1,1,0,1,1] and [1,1,0,1,0] are only compatible for complex

**Table 3:** Discussion of the two arms robot. (Section 11.3).

Case	Conditions	Gröbner basis
Generic case	$l \neq 0,$ $r^2 + z^2 \neq 0,$	$[ 2lc_2 + l^2 - r^2 - z^2 + 1, l^4 - 2r^2l^2 - 2z^2l^2 - 2l^2$ $+ 4l^2s_2^2 - 2r^2 + r^4 - 2z^2 + 2r^2z^2 + z^4 + 1,$ $rl^2 - r - r^3 - z^2r + (2r^2 + 2z^2)c_1 - 2lzs_2,$ $zl^2 - zr^2 - z^3 - z + (2r^2 + 2z^2)s_1 + 2rls_2 ]$
Singular cases		
[1,1,0,1,1]	$l \neq 0, r \neq 0,$ $r^2 + z^2 = 0, z \neq 0$ $l^2 - 1 \neq 0$	$[ 2lc_2 + l^2 + 1, 2zls_2 + r - rl^2,$ $4r(l^2 - 1)c_1 + l^4 + 1 - 4z^2 - 2l^2, -4z^4 + z^2 + 2r^2$ $+ 4zr^2(l^2 - 1)s_1 + l^4z^2 + 2r^2l^4 - 4r^2l^2 - 2l^2z^2 ]$
[1,1,0,1,0]	$l \neq 0, r \neq 0,$ $r^2 + z^2 = 0, z \neq 0$ $l^2 - 1 = 0$	[ 1 ]
[1,0,0,1]	$l \neq 0, r = 0,$ $z = 0, l^2 - 1 \neq 0$	[ 1 ]
[1,0,0,0]	$l \neq 0, r = 0, z = 0,$ $l^2 - 1 = 0$	[ $lc_2 + 1, s_2, c_1^2 + s_1^2 - 1$ ]
[0,1]	$l = 0, r^2 + z^2 - 1 \neq 0$	[ 1 ]
[0,0]	$l = 0, r^2 + z^2 - 1 = 0$	[ $c_2^2 + s_2^2 - 1, c_1 - r, s_1 - z$ ].

values of the parameters ( $r = \pm iz \neq 0$ ) and have no interest for the real configuration.

For the cases [1,0,0,1] and [1,0,0,0] the end of the robot is at the origin. For [1,0,0,1] the system is inconsistent as the length of the second arm is different from  $\pm 1$ . For [1,0,0,0] the angle  $\theta_1$  is free whether  $\theta_2$  is  $\pi$  for  $l = 1$  or 0 for  $l = -1$ , corresponding to the same physical solution.

The cases [0,1] and [0,0] correspond to the degenerate robot of length  $l = 0$ . For [0,1] the system is inconsistent as the position of the end of the robot arm is not at distance 1 of the origin. For [0,0] the angle  $\theta_1$  is determined, whether  $\theta_2$  is free, corresponding to the degeneration of the configuration.

GENCASE reveals itself to be very useful for this problem.

#### 11.4. Load-flow problem for a 3 nodes Electrical Network

In Montes (1995, 1998) we studied the load-flow problem for electrical networks. The equations are polynomial systems with parameters. The following system concerning a concrete three nodes electrical network is considered:

$$\begin{aligned}
14 - 12e_2 - 110f_2 - 2e_3 - 10f_3 - P_1 &= 0, \\
2397 - 2200e_2 + 240f_2 - 200e_3 + 40f_3 - 20Q_1 &= 0, \\
16e_2^2 - 4e_2e_3 - 20e_2f_3 + 20e_3f_2 + 16f_2^2 - 4f_2f_3 - 12e_2 + 110f_2 - P_2 &= 0, \\
2599e_2^2 - 400e_2e_3 + 80e_2f_3 - 80e_3f_2 + 2599f_2^2 - 400f_2f_3 \\
- 2200e_2 - 240f_2 - 20Q_2 &= 0.
\end{aligned}$$

Here  $e_2, f_2, e_3, f_3$  are the variables, representing real and imaginary parts of voltages, and  $P_1, Q_1, P_2, Q_2$  are the parameters representing the real and imaginary components of powers. We use the orders  $\text{lex}(e_2, f_2, e_3, f_3)$  and  $\text{lex}(P_1, Q_1, P_2, Q_2)$ . As a result of the application of DISPGB the following polynomials appear:

$$\begin{aligned}
h_1 &= P_1 - 20 \\
h_2 &= 20 Q_1 - 3497, \\
h_3 &= 6999 P_2 - 800 Q_2 \\
g_1 &= 7786876 h_1 - 79955 h_2, \\
g_2 &= 400 h_1^2 + h_2^2, \\
g_3 &= 7995500 h_1 + 1946719 h_2 \\
g_4 &= h_2 Q_2 (1814947407168 h_2 + 20 778249588225 h_1) + 1392556035 h_3^2.
\end{aligned}$$

The discussion provided by DISPGB and GENCASE is given in table 4.

**Table 4:** Discussion of the load-flow problem

Case	Conditions	Leading power products
[1,1] = Generic case	$g_1 \neq 0, g_2 \neq 0$	$f_3^2, e_3, f_2, e_2$
Special cases		
[1,0,1]	$g_1 \neq 0, g_2 = 0, g_3 h_3 \neq 0$	$f_3, e_3, f_2, e_2$
[1,0,0,1]	$g_1 \neq 0, g_2 = 0, g_3 h_3 = 0, g_4 \neq 0$	1
[1,0,0,0]	$g_1 \neq 0, g_2 = 0, g_3 h_3 = 0, g_4 = 0$	$e_3, f_2, e_2$
[0,1]	$g_1 = 0, h_2 \neq 0$	$f_3, e_3^2, f_2, e_2$
[0,0,1]	$g_1 = 0, h_2 = 0, h_3 \neq 0$	1
[0,0,0]	$g_1 = 0, h_2 = 0, h_3 = 0$	$e_3^2, f_2, e_2$

In this example, DISPGB gives directly the generic case, and GENCASE does not need to improve the result. We only give the sets of leading power products of the different bases, in order to illustrate how the initial goal has been reached (the bases themselves have large coefficients and would confuse the results). Except for the inconsistency cases [1,0,0,1] and [0,0,1], all other have distinct sets of leading power products and cannot be reduced. Obviously the minimal singular variety in this example is  $\mathbb{V}(g_1, g_2)$ .<sup>††</sup>

### 11.5. Application to Example 5.1

We give here the complete discussion for the example 5.1 (see table 5) given by DISPGB, where

$$\Delta = (a - 1)(3a - 7) = 3a^2 - 10a + 7.$$

and  $G$  is the basis of the generic case given there.

<sup>††</sup>In the tutorial, the analysis on how the conditions can be used to simplify the output is described.

**Table 5:** Discussion of example 5.1

Case	Conditions	Gröbner basis
[1] = Generic case	$\Delta \neq 0$	$G$
Special cases	$\Delta = 0$	
[0,1]	$\Delta = 0,$ $ba - 8a - 28 + 35b \neq 0$	[ 1 ]
[0,0]	$\Delta = 0,$ $ba - 8a - 28 + 35b = 0$	[ $-696 + 339b + (-176b + 148)u + 252z,$ $-75 + 33b + 63y + (-32b + 67)u,$ $-309b + 204 + 252x + (208b - 152)u$ ]

Finally, we give the time evaluation of applying DISPGB to the examples described in the paper (and two other from the tutorial), using a PC with a Pentium(r) II, Intel MMX(TM) tech., 128 Mb.

Example	Time(sec)	Example	Time(sec)
11.1	8.8	5.1	8.4
11.2	5.2	Sing. points cubic	1.8
11.3	115.9	$2 \times 2$ gen. lin. sys.	8.2
11.4	33.0		

## Acknowledgements

I am very grateful to Volker Weispfenning for his remarks about the convenience of using his algorithm for the load-flow problem (Montes, 1995), for the fruitful discussions and suggestions and for his excellent and hard review work. His comments induced the present work and helped me in the design. I would extend the gratitude to the anonymous referee's of the paper. I am also grateful to his student Michael Pesch for his patience and interest in the use of his software.

I am also highly indebted to Rafael Sendra for his critical and detailed lecture of the preprint and accurate comments that I have carefully incorporated.

## References

- W.W. Adams, A.K. Boyle, (1992). Some Results on Gröbner Bases over Commutative Rings, *Jour. Symb. Comp.*, **13**, 473–484.
- A. Assi, (1994). On Flatness of Generic Projections, *Jour. Symb. Comp.*, **18**, 447–462.
- D. Bayer, M. Stillman, (1987). A theorem on refining division orders by the reverse lexicographic order, *Duke J. Math.*, **55**, 321–328.
- T. Becker, V. Weispfenning, *Gröbner Bases. A Computational Approach to Commutative Algebra*, Springer, New-York, (1993).

- T. Becker, (1994). On Gröbner bases under specialization, *Appl. Algebra Engrg. Comm. Comput.*, **5**, num. 1, 1–8.
- B. Buchberger, “Ein Algorithmus zum Auffinden der Basiselemente des Restklassenringes nach einem nulldimensionalen Polynomialideal”, *Ph.D. Dissertation*, Univ. Innsbruck, Austria, (1965).
- B. Buchberger, “Gröbner bases: An algorithmic method in polynomial ideal theory”, *Recent Trends in Multidimensional System Theory, chap. 6*, N.K. Bose, Ed. Reidel Publ. Comp., (1985).
- D. Cox, J. Little, D. O’Shea, *Ideals, Varieties and Algorithms*, Springer, New-York, (1992).
- D. Duval, (1995). Évaluation dynamique et clôture algébrique en Axiom, *J. Pure Appl. Algebra*, **99**, num. 3, 267–295.
- R. Gebauer, H.M. Möller, *On an installation of Buchberger’s Algorithm*, Springer, New-York, (1987).
- P. Gianni, T. Mora, “Algebraic Solutions of Systems of Polynomial Equations using Gröbner Bases”, *Applied algebra, algebraic algorithms and error-correcting Codes*, Menorca, (1987). *Lect. Notes Comp. Sci.*, **356**, 247–257. Springer, (1989).
- P. Gianni, (1987). “Properties of Gröbner Bases under Specializations”, *EUROCAL’87, Lect. Notes Comp. Sci.*, J.H. Davenport, **378**, 293–297. Springer,
- H.G. Gräb, (1993). On Lucky Primes, *Jour. Symb. Comp.*, **15**, 199–209.
- M. Kalkbrenner, “Solving systems of algebraic equations by using Gröbner Bases”, *EUROCAL’87, European Conference on Computer Algebra.*, *Lect. Notes Comp. Sci.*, J.H. Davenport, ed., **378**, 282–292. Springer, (1987).
- M. Kalkbrenner, (1997). On the stability of Gröbner Bases under specializations, *Jour. Symb. Comp.*, **24**, num. 1, 51–58.
- H.M. Möller, (1988). On the construction of Gröbner Bases using syzygies, *Jour. Symb. Comp.*, **6**, 345–359.
- A. Montes, (1995). Solving the load flow problem using Gröbner bases, *SIGSAM Bull.*, **29**, 1–13.
- A. Montes, (1998). Algebraic solution of the load-flow problem for a 4-nodes electrical network, *Math. and Comp. in Simul.*, **45**, 163–174.
- A. Montes, “Basic algorithm for Specialization in Gröbner basis”, *Actas de EACA-99*, I. Bermejo, Universidad de La Laguna, Tenerife, 215–228. (1999).

- W.Y. Sit, (1992). An Algorithm for Solving Parametric Linear Systems, *Jour. Symb. Comp.*, **13**, 353–394.
- F. Pauer, (1992). On Lucky Ideals for Gröbner Basis Computations, *Jour. Symb. Comp.*, **14**, 471–482.
- V. Weispfenning, (1992). Comprehensive Gröbner Bases, *Jour. Symb. Comp.*, **14**, 1–29.